# Directions for use Read carefully!



# **ECOSOL G10** (Solution for Intravenous Infusion)

## Symptoms and treatment for overdosage

If hyperglycaemia or glucosuria occur, the dosage and infusion rate should be reduced, or an insulin administration is recommended.

# 555 Side effects

Hyperglycemia and renal loss may occur in case of reduced glucose tolerance. These manifestations are normally prevented by reducing the dosage and/or by giving insulin. Enhanced bilirubin and lactate levels may be found if the recommended dosage is exceeded. Glucose solutions are acidic (pH 3.5 - 5.0) and may produce thrombophlebitis.

## Contraindications

Diabetes mellitus (with the exception of hypoglycaemia conditions) Glucose intolerance Hypotonic dehydration if lacking electrolytes are not replaced Overhydration Hypokalaemia

# Drug interaction

The compatibility of any additives to this solution should be checked before use.

**Storage** Do not store above 30°C.

**Presentation** Plastic bottle 500 ml

Reg no.: DKI1277201749B1

Harus dengan resep dokter

# **B BRAUN**



# schwarz

Dokument = 148 x 210 mm DIN A 5 - 1 Seite

10% Glucose Intravenous Infusion BP (Ecosol G10) – EP 19/12609218/0512 – ID Standort Penang

Lätus 7055

G 090306

#### Composition

Each 100 ml solution contains:

| Glucose Mono   | 11.0 g         |             |
|----------------|----------------|-------------|
| Water for Inje | 100 ml         |             |
| Caloric Value  | kJ/l<br>kcal/l | 1700<br>400 |

|                   |            | kcal/l |
|-------------------|------------|--------|
| Osmolarity mOsm/l | Osmolarity | m0sm/l |

# Mechanism of action

Glucose is absorbed from the gastrointestinal tract, and oxidized as a source of energy, or stored in the liver as glycogen. It is the only energy substrate which is directly, instantly and universally utilized by the body. Glucose is vital for the myocardium, brain and nerves.

#### Indications

Energy supply Hypertonic dehydration Vehicle solution for supplementary medication

#### Posology

The maximum dosage is 500 mg/kg/hr (5 ml/kg/hr of Glucose 10 %) over period of less than 24 hours.

Route of administration: I.V.

## Precautions/Warning

Blood glucose, serum electrolytes and water balance should be monitored regularly.

Electrolytes are to be supplemented as required. The compatibility of any additives to these solutions should be checked before use. Glucose solution should not be administered through the same infusion set through which also blood has or may be given because of the risk pseudoagglutination.

#### Cautions

The compatibility of any additives to this solution should be checked before use.

