

## **NAME OF THE MEDICINAL PRODUCT**

Glucose B. Braun 500 mg/ml Solution for infusion

## **COMPOSITION**

1000 ml of solution for infusion contain:

Glucose monohydrate	550.0 g
(equivalent to glucose)	(500.0 g)

### *Excipients:*

Water for injection, hydrochloric acid (for pH adjustment)

## **THERAPEUTIC INDICATIONS**

Administration of glucose for caloric support; carbohydrate component in parenteral nutrition, in particular in high-caloric nutrition regimen or in cases where restricted fluid intake is indicated; treatment of hypoglycemia.

## **CONTRAINDICATIONS**

Hyperglycemia, not responding to insulin doses of up to 6 units insulin/hour, delirium tremens if such patients are already dehydrated, acute states of shock and collapse, metabolic acidosis, hyperhydration, pulmonary edema, acute cardiac failure.

## **UNDESIRABLE EFFECTS**

Undesirable effects are listed according to their frequencies as follows:

Not known: (cannot be estimated from the available data)

### ***Metabolism and nutrition disorders***

Not known: Hospital-acquired hyponatremia

### ***Nervous system disorders***

Not known: Hyponatremic encephalopathy

Hospital-acquired hyponatremia may cause irreversible brain injury and death due to the development of acute hyponatremic encephalopathy.

## **WARNINGS**

Keep out of the sight and reach of children. For single use only. Discard unused contents. Do not reconnect partially used containers. Use only if solution is clear and colourless or slightly yellowish and container and closure are undamaged.

## **NOTE**

*Prescription only*

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

**B. Braun Melsungen AG, 34212 Melsungen, Germany, 01/2018**